

Ceftiofur Sodium for Injection

(Bio Agri Mix)

Ingredient(s):

benzyl alcohol, ceftiofur sodium, water for injection

Use:

Bovine respiratory disease (*Mannheimia haemolytica*, *Pasteurella multocida*, and/or *Histophilus somni*), treatment
Footrot [*Fusiformis necrophorus* (*Fusobacterium necrophorum*) and/or *Porphyromonas levii* (*Bacteroides* spp)], control and/or treatment

Species:

Beef Cattle

RoA:

I.M.

Dose:

1 mL/50 kg (1 mg/kg) b.w. Repeat q24h for total of 3 treatments. Additional treatments may be given on days 4 and 5 for animals which have not recovered after initial 3 injections.

Withdrawal Time:

Meat: 0d

Use:

Bovine respiratory disease (*Mannheimia haemolytica*, *Pasteurella multocida*, and/or *Histophilus somni*), treatment
Footrot [*Fusiformis necrophorus* (*Fusobacterium necrophorum*) and/or *Porphyromonas levii* (*Bacteroides* spp)], control and/or treatment

Species:

Dairy Cattle

RoA:

I.M.

Dose:

1 mL/50 kg (1 mg/kg) b.w. Repeat q24h for total of 3 treatments. Additional treatments may be given on days 4 and 5 for animals which have not recovered after initial 3 injections.

Withdrawal Time:

Meat: 0d Â Milk: 0d

Use:

Urinary tract infections, treatment

Species:

Dogs

RoA:

S.C.

Dose:

0.2 mL/5 kg (2 mg/kg) b.w. Repeat q24h for 5-14d.

Use:

Respiratory disease (*Streptococcus zooepidemicus*), systemic treatment

Species:

Horses

RoA:

I.M.

Dose:

2 mL/50 kg (2 mg/kg) b.w. Repeat q24h until 48h after symptoms have disappeared. If no response is observed within 4-5d, re-evaluate diagnosis.

WT Notes:

This drug is not to be administered to horses that are to be slaughtered for use in food.

Use:

Respiratory disease (bacterial), control/treatment

Respiratory disease, ovine (*Mannheimia haemolytica*), treatment

Species:

Sheep

RoA:

I.M.

Dose:

Lambs: 2 mL/50 kg b.w. (2 mg/kg). Repeat q24h for a total of 3 treatments. Additional treatments may be given on days 4 and 5 for animals which have not recovered after initial 3 injections.

Withdrawal Time:

Meat: 24h

Use:

Respiratory disease (bacterial), control/treatment

Species:

Swine

RoA:

I.M.

Dose:

1 mL/17 kg (3 mg/kg) b.w. Repeat q24h for 3 treatments.

Withdrawal Time:

Meat: 24h