

Chlor 50 Chlortetracycline Premix

(Bio Agri Mix)

Ingredient(s):

chlortetracycline

Use:

Enteritis (bacterial), prevention

Species:

Beef Cattle, Dairy Cattle

RoA:

In the feed

Dose:

Calves up to 136 kg: Feed continuously during periods of early growth.

In the feed: Add 0.5 kg/tonne [0.0055% (55 mg/kg)].

Milk replacer: Add 0.5 kg/tonne [0.0055% (55 mg/kg)].

Withdrawal Time:

Meat: 5d

WT Notes:

Do not administer to lactating dairy cattle.

Use:

Footrot, prevention

Species:

Beef Cattle, Dairy Cattle

RoA:

In the feed

Dose:

Add 200 g/tonne (0.22 mg/kg/day b.w.) or add 640 g/tonne supplement and feed 1 kg/head/day (70 mg/head/day). Feed continuously during exposure period.

Withdrawal Time:

Meat: 5d

WT Notes:

Do not administer to lactating dairy cattle.

Use:

Enterotoxemia (*Clostridium perfringens*), reduction of losses

Species:

Sheep

RoA:

In the feed

Dose:

Feedlot lambs: 0.0022% (22 mg/kg) complete feed. Feed continuously as sole ration while lambs are in feedlot.

Withdrawal Time:

Meat: 4d

Use:

Enteritis (bacterial), treatment

Species:

Swine

RoA:

In the feed

Dose:

Add 1 kg/tonne [0.011% (110 mg/kg)]. Feed continuously as sole ration following the appearance of symptoms and until 3d after symptoms disappear. Prestarter feeds and feeds to be fed for up to and including 6 weeks of age are required to contain 0.022%.

Withdrawal Time:

Meat: 7d

Use:

Enteritis (bacterial), prevention

Species:

Swine

RoA:

In the feed

Dose:

0.5 kg/tonne [0.0055% (55 mg/kg)]. Feed continuously as sole ration during early growth period up to 32 kg (70 lb) b.w. Prestarter feeds and feeds to be fed for up to and including 6 weeks of age are required to contain 0.022%.

Withdrawal Time:

Meat: 7d

Use:

Hexamitiasis, control/prevention

Species:

Turkeys

RoA:

In the feed

Dose:

0.0055% (55 mg/kg) complete feed: Feed as sole ration for as long as birds are expected to be exposed to infection.

Withdrawal Time:

Meat: 7

Use:

Hexamitiasis, treatment

Species:

Turkeys

RoA:

In the feed

Dose:

0.011% (110 mg/kg) complete feed: Feed as sole ration when symptoms appear and continue for a few days after symptoms disappear.

Withdrawal Time:

Meat: 7d

Use:

Synovitis (infectious - *Mycoplasma synoviae*), treatment

Species:

Turkeys

RoA:

In the feed

Dose:

0.022% (220 mg/kg) complete feed: Feed as sole ration when symptoms appear and continue for at least 2 weeks after symptoms disappear.

Withdrawal Time:

Meat: 7d