

Tylosin Soluble Powder

(Bio Agri Mix)

Ingredient(s):

tylosin tartrate

Use:

Necrotic enteritis (*Clostridium perfringens*), treatment

Species:

Chickens

RoA:

dw

Dose:

Broiler chickens: 100-150 mg/L, depending upon severity of infection for 5d.

100 g: To make 0.4-0.6 g/4 L, mix 100 g/1,000 L water.

500 g: To make 0.4-0.6 g/4 L, mix 500 g/5,000 L water.

Prepare fresh solution q3d.

Withdrawal Time:

Meat: 0d

WT Notes:

Do not use in laying birds.

Use:

Chronic respiratory disease/infectious sinusitis, control/prevention

Chronic respiratory disease/infectious sinusitis, treatment

Species:

Chickens

RoA:

dw

Dose:

Prevention at time of vaccination or other stress: 2 g/4 L (500 mg/L) for 3d. Repeat for 1d at 3-4 weeks of age to coincide with vaccination or other stress.

Treatment: 2 g/4 L (500 mg/L) for 1-5d depending on severity of infection.

Prepare fresh solution every 3d.

100 g package: To make 2 g/4 L solution, mix 100 g/200 L water.

500 g package: To make 2 g/4 L solution, mix 500 g/1,000 L water.

Withdrawal Time:

Meat: 1d

WT Notes:

Do not use in laying birds.

Use:

Porcine proliferative enteritis (ileitis), treatment

Species:

Swine

RoA:

dw

Dose:

1 g/12 L drinking water for 14d, or 1 g/12L for 7d followed by 110 mg/kg tylosin phosphate premix in the feed for 7d.

100 g package: To make 1 g/12 L solution, mix 100 g/1,200 L water.

500 g package: To make 1 g/12 L solution, mix 500 g/6,000 L water.

Withdrawal Time:

Meat: 0d

WT Notes:

Treated pigs administered 1 g of tylosin per 4 L (250 mg/litre) of drinking water must not be slaughtered for use in food for at least 48 hours after the latest treatment with this drug.

Use:

Swine dysentery, treatment

Species:

Swine

RoA:

dw

Dose:

1 g/4 L water for 3-10d.

100 g package: To make 1 g/4 L solution, mix 100 g/400 L water.

500 g package: To make 1 g/4 L solution, mix 500 g/2,000 L water.

Withdrawal Time:

Meat: 2d

WT Notes:

Treated pigs administered 1 g of tylosin per 4 L (250 mg/litre) of drinking water must not be slaughtered for use in food for at least 48 hours after the latest treatment with this drug.

Use:

Chronic respiratory disease/infectious sinusitis, control/prevention
Chronic respiratory disease/infectious sinusitis, treatment

Species:

Turkeys

RoA:

dw

Dose:

Prevention: 2 g/4 L (500 mg/L) for 5d. Repeat for 2d at 4 weeks of age.

Treatment: 2 g/4 L (500 mg/L) for 3d. Depending on severity of infection treatment can be given for 2-5d.

Prepare fresh solution every 3d.

100 g package: To make 2 g/4 L solution, mix 100 g/200 L water.

500 g package: To make 2 g/4 L solution, mix 500 g/1,000 L water.

Withdrawal Time:

Meat: 3d

WT Notes:

Do not use in laying birds.